

Sedimentation Across the Tidal-Fluvial Transition in the Lower Fraser River, Canada

Shahin E. Dashtgard^{1*}, Jeremy G. Venditti², Philip R. Hill³, Chad F. Sisulak¹, Stacy M. Johnson¹, and Andrew D. La Croix¹

¹Applied Research in Ichnology and Sedimentology (ARISE) Group, Department of Earth Sciences, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6

*email: sdashtgard@gmail.com

²Department of Geography, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6

³Geological Survey of Canada, Natural Resources Canada, Pacific Geosciences Center, Sydney, British Columbia V8L 4B2

doi: 10.2110/sedred.2012.4.4

ABSTRACT

The Fraser River is the largest undammed river on the west coast of North America. In its lower reaches, a saltwater wedge intrudes up to 30 km inland during mixed semi-diurnal tidal cycles that range up to 5.3 m in height. Sediments deposited in the lower reaches of the Fraser River show distinctive characteristics that reflect the relative control of river versus tidal processes, as well as the persistence of saline water at each point along the channel.

Grain-size trends along the river are controlled by the hydrodynamics in each distributary. Mud deposition is concentrated in the zone of saltwater-freshwater mixing. Coarse-grained sand and mud/fine-grained sand deposition is largely seasonally controlled, wherein bed material (diameter > 0.177 mm) is deposited during the waning freshet, and washload transported mud and fine-grained sand (< 0.177 mm) is deposited during the late-stage waning freshet flow and during base flow.

The diversity and density of bioturbation changes according to the volume and residence time of brackish water at the bed. Higher salinity water and greater durations that saline water is sustained at any locale, supports a more diverse and uniformly distributed trace assemblage. With decreasing salinity, the trace assemblage decreases in diversity and bioturbation becomes more sporadically distributed. This results in a reduction of infaunal diversity from 100% on the nearly fully marine tidal flats in the abandoned part of the lower delta plain, to 14% in intertidal sediments of the brackish-water reach.

The character of the sediments deposited across the tidal-fluvial transition provide criteria for differentiating sediments deposited within freshwater-tidal reaches, brackish-water-tidal reaches, and mixed tidal-fluvial distributaries. These data are presented as a process-based analog for tidal-fluvial sediments preserved in the rock record. The results can be used to predict changes in facies character across the tidal-fluvial transition of similar tide-influenced, river-dominated systems or their rock-record equivalents.

INTRODUCTION

With increasing interest in recognizing differences in the sedimentological and ichnological character of sediments deposited across the tidal-fluvial transition, it is necessary to study a variety of modern analogs subjected to varying degrees of fluvial and tidal input. One such analog is the Fraser River, British Columbia, Canada, which in its lower reaches is affected by strong tidal flow. The grain sizes of sediments

deposited in the channel are strongly linked to the degree of tidal flow relative to river flow, and to the extent and duration of saltwater intrusion at sites of sediment deposition. Consequently, the sedimentological and ichnological character of the sediments reflects the depositional conditions under which they were deposited, and hence, can be used as a proxy for establishing the volume and duration of saline water intrusion and the relative input of tidal versus river flow.

Understanding sediment deposition across the tidal-fluvial transition has received increasing attention in recent years (e.g., Dalrymple and Choi 2007) in large part because subsurface reservoirs of the bitumen-hosting middle McMurray Formation in northeast Alberta are interpreted as paleo-estuarine channel bars (Pemberton and Wightman 1992; Ranger and Pemberton 1997; Musial et al. 2012). The lateral and vertical extent of mud beds and muddy bedsets interbedded with sand in the channel bars is the dominant control on reservoir compartmentalization, and can strongly impact the economic viability of exploiting hydrocarbons contained in these deposits (Strobl et al. 1997; Hein and Cotterill 2006). To better understand mud distribution in tide-influenced river channels, a range of modern analogues are being evaluated across the spectrum from tide-dominated (Dalrymple et al. 2003; Choi et al. 2004; Pearson and Gingras 2006), to fluvially influenced, tide-dominated (Smith 1988; Gingras et al. 1999; Smith et al. 2009), to mixed tidal-fluvial (Johnson 2012), to tide-influenced, fluvially dominated (Sisulak and Dashtgard 2012) to fluvially dominated (Smith et al. 2009; Smith et al. 2011) settings.

The lower Fraser River (Fig. 1) is a natural laboratory for studying sediment deposition in tide-influenced, fluvially dominated channels and mixed tidal-fluvial channels. The availability of historical and real-time hydraulic data (e.g., Johnson 1921; Ages 1979; Thomson 1981), the fact that the Fraser River is undammed along its entire length, and the extensive studies of sediment transport in the river (e.g., Kostaschuk and Luternauer 1989; Church and Krishnappan 1995; Kostaschuk and Villard 1996; Kostaschuk et al. 1998; McLean et al. 1999; Kostaschuk and Best 2005) provide datasets that can be used to link sedimentological, ichnological, and architectural characteristics of the sediments to the hydrodynamic and chemical conditions under which they were deposited and colonized by infauna.

THE FRASER RIVER

The Fraser River drains 228000 km² of mountainous terrain and has a mean annual river discharge of 2710 m³s⁻¹ at Hope, BC, ~165 river km upstream of the delta front. Low flow rates in winter are ~1000 m³s⁻¹, and the annual peak flow ranges between 5130 and 15200 m³s⁻¹ with a mean of 8642 m³s⁻¹ (Water Survey of Canada, <http://www.wateroffice.ec.gc.ca/>, accessed Sept 7, 2012). At low flow, the saltwater wedge (SWW) intrudes ~30 km upriver (Kostaschuk and Atwood 1990). During high river flows,

the SWW is displaced towards the mouth of the channel. At river flows between 5000 and 7000 $\text{m}^3 \text{s}^{-1}$ (at Hope), a SWW has been observed to penetrate only a few kilometers up the Main Channel (red line, Fig. 1) during spring high tides (Milliman 1980). At high peak flows ($> 8000 \text{ m}^3 \text{ s}^{-1}$ at Hope), however, the SWW likely does not enter the Main Channel.

In its tide-influenced lower reach, the Fraser River bifurcates into two distributaries, the North Arm and the Main Channel (Fig. 1). The North Arm bifurcates again to form the Middle Arm. Observations made in 1970 at a flow rate close to the mean annual peak rate (8670 $\text{m}^3 \text{ s}^{-1}$) indicated that the North and Middle arms carried $\sim 7\%$ and $\sim 5\%$ of channel flow respectively (WCHL 1977). The Main Channel carried 70% of the flow and its distributary, Canoe Pass, carried 18% of the flow (Fig. 1).

Sediment flux in the Fraser is washload dominated. Washload is sediment not present in the channel bed and lower banks of the river, and that stays in suspension through a defined river reach (Church 2006). In the Main Channel and the North Arm, the washload is composed of fine sand ($< 0.177 \text{ mm}$), silt, and clay, whereas in the Middle Arm and probably Canoe Pass, similar grain sizes are deposited on the channel bed during non-peak flows (Johnson 2012). Landward of New Westminster, before the Fraser bifurcates into its various distributaries, McLean et al. (1999) reported that between 1966 and 1986, the total sediment load was $17 \times 10^6 \text{ t yr}^{-1}$. Forty-nine percent of the sediment load was silt-

sized material, whereas clay made up 15%. Sand-sized sediment constituted 36% of the total load, roughly half of which was carried as washload into the lower river reaches. The remaining 18% of the sand volume was carried as bed-material load, defined as sediment present in the bed or lower banks of the river and transported as bedload or in temporary suspension (Church 2006). The distribution of the sediment load through the various distributaries and delta is, however, only understood conceptually from estimates of aggradation-degradation of the channel and from sediment deposits.

Anthropogenic modification of the lower Fraser River and delta is significant. The channel banks were first diked between 1913 and 1919 (Johnson 1921) essentially stopping lateral and downstream migration of distributaries. The dike system has been gradually expanded to restrict 90% of the river in the Fraser Valley, downstream of Hope (Fig. 1). Extensive placer mining (Nelson and Church 2012) along the Fraser River over the past 150 years has contributed to periods of high sediment yield, which have influenced the delta topography. For example, the South Arm Marshes bar complex (discussed later) is thought to have aggraded significantly following the influx of sediment from placer mining activities in the mid to late 1800s (Public-Works-Canada 1957; Hales 2000).

The Main Channel has been regularly dredged since 1945 to provide a deep water navigation channel up to New Westminster. Prior to that, the sandy bed-material load passed through the estuary, and $3.0 \times 10^6 \text{ t yr}^{-1}$ of sand was

delivered to the delta front. Over-deepening of the channel due to dredging has produced a knick-point that has been migrating upstream of New Westminster towards Mission, lowering the bed elevations and recruiting an additional $1.05 \times 10^6 \text{ t yr}^{-1}$ of sand. As such, $\sim 4.25 \times 10^6 \text{ t yr}^{-1}$ of sand are now delivered to the river of which 70% is dredged. Over the past 50 years, dredging has removed an average of $2.95 \times 10^6 \text{ t yr}^{-1}$ of bed-material load (sand $> 0.177 \text{ mm}$; Church 2010), which restricts the fluvial sand supply to the delta front to $\sim 1.3 \times 10^6 \text{ t yr}^{-1}$.

Regardless of the anthropogenic modification of the Fraser River and delta, the Fraser is undammed and only one significant tributary has regulated flow: a rarity amongst major rivers. As such, precipitation and climatic conditions in the mountainous catchment basin are reflected in the depositional conditions in the lower reaches. Seasonal variations in river discharge, and brackish water and tidal inundation up the Fraser distributaries reflect natural hydraulic conditions. This makes the Fraser an excellent analog for comparing sedimentological and ichnological trends to hydrodynamic processes and water salinity.

SEDIMENTOLOGICAL TRENDS IN THE LOWER FRASER RIVER

There are pronounced sedimentological trends in the lower Fraser River related to: 1) channel morphology and grain size, 2) mud content and distribution, 3) bedding cyclicity and architecture, and 4) sedimentary structures.

1) Perhaps the most obvious trends in the lower Fraser River are the pronounced changes in grain size and channel morphology between the non-tidal freshwater reach, the freshwater-tidal reach, and the brackish water-tidal reach (Fig. 1). The Fraser experiences tides $\sim 90 \text{ km}$ inland to Mission, but only the most seaward 30 km is regularly inundated by brackish water. The remaining 60 km of tide-influenced river is freshwater and tidal. The Fraser River exhibits a major grain size and morphological change 15 km upstream of Mission where the channel changes from a wandering gravel-bed river (Church 1983; McLean 1990; Ham 2005) to a sinuous sand-bed channel (Venditti et al. 2010) with in-channel (tidal) bars. The position of this transition is thought to be topographically controlled (Venditti et al. 2010), although tides may play a secondary role. The wandering gravel-bed river occurs wholly in the freshwater, non-tidal reach, whereas the sand-bed channel extends through the freshwater-tidal and brackish water-tidal reaches.

In the brackish water-tidal reach, saltwater intrusion (brackish-water + tidal, Fig. 1) is periodic (depending on river discharge) and the tidally affected portions of the bars contain significant volumes of both sand and mud (Fig. 2). The channels in this reach also show evidence of lateral and downstream migration (prior to the channel being diked) and tend to




Figure 1: Composite airphoto of the lower Fraser River from Hope, British Columbia, Canada to the delta front (Image source: Google Earth). At the top of the image, the general hydrodynamic and salinity conditions (e.g., sustained brackish water + tidal) are defined, as is the general channel morphology (e.g., straight) and dominant grain size(s) (e.g., mixed sand-mud). The approximate position of the defined zones are bracketed by the solid (applicable to all distributaries) and dashed (varies between distributaries) black lines above the airphoto. The red lines mark the maximum upstream extent of saltwater intrusion during high river flow (freshet) and the yellow line marks the maximum upstream intrusion of saltwater during base flow. The orange line is the approximate upstream location of observable tides. The percentages on the left hand side of the photo are the estimated percent volume of flow that is transported down each distributary (WCHL, 1977). The thick black line on the airphoto marks the landward boundary of the Fraser delta. The location of the Fraser River delta in Canada is shown in the inset map. The blue circles mark the position of cities and towns along the lower Fraser River that are referred to in the text and in Figure 2.




Figure 2: Downstream change in the grain-size fraction in the Fraser River from Hope to Sand Heads (just seaward of the break in slope between the lower delta plain and the delta front). Each data point on the graph averages the results of samples collected from the channel base to the channel margins. Data compiled from McLaren and Ren (1995), Venditti (unpublished data), Venditti et al. (2010), McLean (1990), and Ham (2005). Modified from Venditti et al. (2010). DF = delta front, New West = New Westminster.

be straighter than in the freshwater-tidal reach. Finally, in the river reaches that experience more sustained brackish water, the channels are straight and the sediments in this region are mixed sand and mud (Figs. 1 & 2).

2) The most significant change observed in the sedimentology between the freshwater-tidal reach and the brackish water-tidal reach is in the distribution of mud. In the freshwater-tidal reach, mud-dominated beds and bedsets are typically thin (< 10 cm thick) and are laterally continuous for meters to tens of meters. The channel bottom and intertidal banks are sand-dominated (Fig. 2), and mud deposited during base flow is typically eroded during the freshet. In the brackish water-tidal reach, mud deposition is increasingly widespread (Fig. 3). Mud beds and mud-dominated bedsets thicken (to > 1 m thick) and are laterally continuous for hundreds of meters to kilometers in the intertidal zone and likely the upper subtidal zone (Johnson 2012; Sisulak and Dashtgard 2012). The thickness and lateral continuity of muddy bedsets is linked, at least in part, to the persistence and duration of saline water and the development of a turbidity maximum, and in part to the relative influence of fluvial versus tidal processes on sediment deposition. This is best expressed in the differences in the mud content on bars in the Main Channel (South Arm Marshes (SAM) bar complex, Fig. 3B; Sisulak and Dashtgard 2012) and in the Middle Arm (Fig. 3C-E; Johnson 2012).

3) The cyclicity and architecture of bedding within bars changes with position along the channel, and reflects both the relative influence of fluvial and tidal processes, and the variability in the amount of mud deposited in and along the margins of the river. This, in turn, is linked to the position of the saltwater wedge. Johnson (2012) found significant uniformity in the thickness of sand beds and sand-mud interbeds (Fig. 4) in the admixed tidal-fluvial (i.e., tidal and fluvial processes equally control sediment deposition) Middle Arm. She attributed this

uniformity in bedding to the relatively strong tidal influence on sediment deposition. In the tide-influenced, fluvially dominated Main Channel, Sisulak and Dashtgard (2012) found that along the margins of the SAM bar complex (Fig. 3B), sand beds were non-uniform in thickness, and sand-mud interbedding was non-rhythmic (Fig. 5). They attributed this relation to the fact that sand transport and deposition was dominated by river flow, which

changes significantly from year to year. Mud beds, however, were deposited during lower flow and base-flow conditions, when estuarine circulation was re-established and the washload started to settle out of suspension. Landward of the maximum upstream position of the saltwater wedge, mud deposition is limited, and the seasonal cyclicity between sand and mud interbeds is not observed. Consequently, vertical bedding in the freshwater-tidal and freshwater-non-tidal reaches (Fig. 1) are either sand-dominated or gravel dominated, with rare thin and discontinuous mud interbeds.

4) The sedimentary structures preserved within bars along the Fraser River record tidal influence on sediment deposition. In the admixed tidal-fluvial Middle Arm, bars contain abundant mm- to cm-scale sand-mud laminasets that occasionally occur in full sets of 12-14 couplets and are interpreted as tidal rhythmites. Tidal rhythmites are still present in bars through the tide-influenced, fluvially dominated part of the Main Channel and Canoe Pass (Fig. 6B), but the number of rhythmites is greatly reduced. Cross-bedding similar to sigmoidal cross-bedding (Kreisa and Moiola 1986) was also observed in sandy dunes developed in the intertidal zone in Canoe Pass (Fig. 6A), although this bedding style is attributed to tidal modification of fluvially formed dunes. Flaser bedding was




Figure 3: A) Percent mud map for the Fraser River distributaries compiled by the Geological Survey of Canada. The location of maximum saltwater intrusion is shown. The dashed black boxes mark the position of Figures 3B-E. B-E) Detailed grain size maps of four bars along the Main Channel and Middle Arm of the Fraser River. Grain size maps are the result of intertidal surface sediment sampling and subaqueous grab sampling. The numbers "4" and "5" shown in white circles on E and B, respectively, indicate the position of the vibracore strip logs in figures 4 and 5. Figure 3B is of the South Arm Marshes (SAM) bar complex, and is modified from Sisulak and Dashtgard (2012). Figures 3C-E are modified from Johnson (2012).

density is indicative of the relative persistence of brackish-water conditions along the channel profile, and can be used as proxy for determining salinity in the channel. The interbedding of bioturbated and non-bioturbated beds is indicative of seasonal changes in river discharge and the corresponding decrease in salinity and increase in the sedimentation rate.

with very rare occurrences of BI 1-2; Fig. 6F) of small diameter *Planolites*- to *Thalassinoides*-like, horizontal burrow networks. For reference, the bioturbation index is a non-linear scale, from 0 to 6, of biogenic reworking of the sediment: the scale is included in the legend in Figure 6.

With increasing saltwater intrusion and the persistence of brackish water conditions at any point along the channel profile, there is a corresponding increase in the diversity and density of bioturbation (Fig. 6B-E; Chapman and Brinkhurst 1981; Johnson 2012; Sisulak and Dashtgard 2012). In bars through the tide-influenced, fluviially dominated Main Channel, the trace density increases from BI 0-2 (Fig. 6D-E) to BI 0-4 (Fig. 6B-C), where the burrow density in individual bedsets is dependent upon both the persistence of saltwater and the sedimentation rate. The trace assemblage comprises a low diversity suite of *Arenicolites*-, *Skolithos*-, and *Polykladichmus*-like burrows (Fig. 6B-E), most of which are the vertical dwellings of polychaetes. In the tidal-fluvial Middle Arm, burrows produced by marine organisms are maintained year-round due to the persistence of brackish water and a reduced sedimentation rate resulting from lower river discharge. As such, Middle Arm sediments tend to exhibit a higher density (BI 2-6) and diversity of burrowing with the addition of both large and small *Siphonichmus*-like (bivalve-generated) traces (Johnson 2012). The traces in the tidal-fluvial Middle Arm are larger on average than those found in the tide-influenced, fluviially dominated Main Channel.

The increase in the density and diversity of traces with increasing brackish-water influence in the channels is relatively subtle when compared to the large increase in trace density and diversity from the channels onto the lower delta plain tidal flats of the Fraser River (Dashtgard 2011b), and to the nearly fully marine tidal flats in Boundary Bay (Fig. 1; Dashtgard 2011a). The brackish water-tidal reach of the Main Channel has an infaunal diversity of 3, which is 14% that of the nearly fully marine tidal flats of Boundary Bay. In the tidal-fluvial Middle Arm, the infaunal diversity of intertidal zone sediments doubles from 3 to 6 common animals. There is a significant increase in infaunal diversity from the intertidal zone of the Middle Arm to the active lower delta plain tidal flats, with an increase in infaunal diversity from 6 to 14 common animals. However, the lower delta plain tidal flats show a diversity reduction of 34% from the Boundary Bay tidal flats (Dashtgard 2011a; Dashtgard 2011b).

The most significant reduction in infaunal diversity occurs from the lower delta-plain tidal flats into the distributaries of the lower Fraser River. The low diversity and density of infauna observed in the channels is distinctive of the brackish-water reaches of the Fraser River, and provides a useful indicator of channel versus non-channel, marginal-marine facies. The upstream decrease in trace diversity and




Figure 5: Strip log for a vibracore taken from the South Arm Marshes bar complex (tide-influenced, fluviially dominated), an in-channel bar along the Main Channel (from Sisulak and Dashtgard, 2012). Note the irregular thickness of mud-dominated and sand-dominated bedsets between 1.25 and 3.0 m depth. The thickness of sand beds is controlled by the river discharge volume, which varies annually. Similarly, mud bed thicknesses are partly controlled by river discharge. The location of this core is shown on Figure 3B. The symbol legend is shown in Figure 4.




Figure 4: Strip log and symbol legend for a vibracore taken from a point bar in the tidal-fluvial Middle Arm, lower Fraser River (modified from Johnson (2012)). Note the regular interbedding and consistent thicknesses of both sand and mud beds. This rhythmicity of bedding is considered to reflect increased tidal control on sediment deposition. The location of this core is shown in Figure 3E. The blue line labeled “MLT” is the mean low tide line.

commonly observed in the intertidal and upper subtidal portions of the Main Channel bars, and mud rip-ups were exceedingly common due to the annual deposition of mud beds and subsequent erosion during the freshet (Sisulak and Dashtgard 2012).

ICHOLOGICAL TRENDS IN THE LOWER FRASER RIVER

The ichnological character of sediments in the lower Fraser River is consistent with the expected trace assemblage for brackish-water settings (e.g., Pemberton and Wightman 1992; MacEachern and Gingras 2007), in that the trace assemblage on all bars in the tide-influenced part of the river is dependent upon the persistence of brackish water at each locale. In the freshwater-non-tidal and freshwater-tidal reaches of the Fraser River, bioturbation is very rare and is limited to sporadically distributed, thin (<2 mm diameter) oligochaete burrow networks. The trace assemblage in these zones is best described as a very low density (bioturbation index (BI) 0

**IMPLICATIONS FOR
SEDIMENT DEPOSITION
ACROSS THE TIDAL-FLUVIAL
TRANSITION**

The lower Fraser River is an excellent analog for studying changes in the sedimentology and ichnology of channel-associated deposits across the tidal-fluvial transition in a tide-influenced, fluvially dominated river. The character of sediments can be linked to the hydrodynamic conditions in the distributaries and to water salinity at various bars along the channel. Based on the results of our research, we make the following observations for the purpose of application to the rock record.

1) Grain-size distributions along the lower Fraser River are linked to tidal influence, and to flow reduction or flow reversal during the flood tide. The river is gravel-bedded upstream of the landward extent tidal effects. The river transitions to a sand-bedded river ~15 km upstream of the freshwater-tidal reach, where tides may play some role in determining this transition. The Main Channel and most of the distributaries are also sand-bedded along the brackish water-tidal reach; however, mud is commonly deposited along the channel margins and in the upper part of in-channel bars in the Main Channel, and on the channel floors and bars of distributaries that carry lower volumes of the river water (e.g., Middle Arm).

2) Mud deposition is concentrated in the zone of freshwater-saltwater mixing (i.e., the turbidity maximum), such that the amount of mud preserved in the bars is linked to the persistence of saltwater at any point along the channel axis. In tidal-fluvial and tide-influenced, fluvially dominated settings, intertidally exposed mud beds are laterally continuous for up to 1 km (Johnson 2012). The lateral continuity of mud beds decreases rapidly landward of the zone of saltwater intrusion.

3) In the brackish-water influenced reaches of the lower Fraser River, coarse sand (> 0.177 mm) is deposited during the waning freshet. Clay, silt and fine sand (< 0.177 mm) are deposited during late-stage waning freshet flow and during base flow, although mud deposition in the Main Channel is restricted to the channel margins. This results in a seasonal cyclicity to sand-mud interbedding (Sisulak and Dashtgard 2012). Seasonal cyclicity is generally not discernable in sediments landward of the brackish-water influenced zone.

4) The diversity of infauna (and their traces) exhibits the greatest decrease from the lower delta plain tidal flats into the channels. In tidal-fluvial and tide-influenced, fluvially dominated channels, infaunal diversity is low: 14-28% of the diversity of the nearly fully marine tidal flats at Boundary Bay. Trace diversity across the tidal-fluvial transition decreases with decreasing saltwater intrusion, suggesting that trace diversity can be used to predict the salinity conditions under which the sediments were colonized.

For tidally influenced rivers, this would correspond to a position along a channel.

5) Trace density is controlled by salinity and the sedimentation rate. Trace density decreases and becomes more sporadically distributed with decreasing salinity, although tidal-fluvial and tide-influenced, fluvially dominated upper bar and channel-margin deposits show BI values of 2-6 and 0-4, respectively. The sedimentation rate also exerts a major control on trace density in that high rates of sediment deposition translate into low density burrowing. In the bars in the tidal-fluvial Middle Arm, this is manifest as lower BI values in sand beds versus mud beds

(Johnson 2012). In the tide-influenced lower reach of the Main Channel, the SAM bar complex, sand beds in the upper part of the bar tend to lack bioturbation or are burrowed from the top down, and mud beds show the highest levels of burrowing (Sisulak and Dashtgard 2012). The correlation between bioturbation and grain size further emphasizes the seasonal cyclicity recorded by interbedded sand and mud beds in brackish-water influenced channels.

Significant variability occurs between the distributaries of the lower Fraser River, such that one style of sedimentation cannot be applied to all channels. However, the sedimentological and




Figure 6: X-radiographs of box cores taken from bars along the lower Fraser River. The position of each box core is shown on the air photo at the bottom of the figure. The two vertical columns on the right-hand side of each x-radiograph show the general lithology and bioturbation intensity (recorded as the bioturbation index). In figures 6B and 6C, the solid horizontal black lines labeled 1, 2 and 3 define the tops of seasonal cycles of deposition. Abbreviations for sedimentary structures (squares with arrows): current ripple lamination (cr); muddy current ripple lamination (mcr); planar lamination (pl); sigmoidal cross bedding (SXB); tidal rhythmite (tr); trough cross-bedding (txb). Abbreviations for ichnological/biogenic structures (circles with arrows): Arenicolites (Ar); bird footprint (fp); Polykladichnus (Pk); Skolithos (Sk); threadworm burrow (t); wood (wd).

ichnological character of the sediments deposited at any location in the river can be used to predict the depositional conditions (processes) under which those sediments were deposited.

ACKNOWLEDGMENTS

The authors wish to thank the people who generously supported this research. SED's research is supported by Nexen Inc., Imperial Oil Ltd., Statoil ASA, Suncor Energy, and the Natural Sciences and Engineering Research Council (NSERC). JGV's research is supported by NSERC. Earth Sciences Sector Publication No. 20120301. Thanks to Kimberley Jenner, Wayne Wright, and Xavier Janson for their reviews of this manuscript.

REFERENCES

- AGES, A., 1979, The salinity intrusion in the Fraser River: salinity, temperature and current observations 1976, 1977, p. 193.
- CHAPMAN, P.M., and BRINKHURST, R.O., 1981, Seasonal changes in interstitial salinities and seasonal movements of subtidal benthic invertebrates in the Fraser River estuary, B.C.: *Estuarine, Coastal and Shelf Science*, v. 12, p. 49-66.
- CHOI, K.S., DALRYMPLE, R.W., CHUN, S.S., and KIM, S.-P., 2004, Sedimentology of Modern, Inclined Heterolithic Stratification (IHS) in the Macrotidal Han River Delta, Korea: *Journal of Sedimentary Research*, v. 74, p. 677-689.
- CHURCH, M., 1983, Pattern of instability in a wandering gravel-bed channel, *in* Collinson, J.D., and Lewin, J., eds., *Modern and Ancient Fluvial Systems*: London, International Association of Sedimentologists, p. 169-180.
- CHURCH, M., 2006, Bed material transport and the morphology of alluvial river channels: *Annual Review of Earth and Planetary Sciences*, v. 34, p. 325-354.
- CHURCH, M.A., 2010, Sustaining Fraser Estuary: Bases for a sediment management plan, *in* Program, F.E.M., ed.: Vancouver, Canada, University of British Columbia, p. 39.
- CHURCH, M.A., and KRISHNAPPAN, B.G., 1995, Sediment sources, transport processes, and modeling approaches for the Fraser River *in* Canada, E., ed., *Fraser River Action Plan*: Vancouver, Canada.
- DALRYMPLE, R.W., and CHOI, K.S., 2007, Morphologic and facies trends through the fluvial-marine transition in tide-dominated depositional systems: A schematic framework for environmental and sequence-stratigraphic interpretation: *Earth -Science Reviews*, v. 81, p. 135-174.
- DALRYMPLE, R.W., BAKER, E.K., HARRIS, P.T., and HUGHES, M.G., 2003, Sedimentology and stratigraphy of a tide-dominated, foreland-basin delta (Fly River, Papua New Guinea), *in* Sidi, F.H., Nummedal, D., Imbert, P., Darman, H., and Posamentier, H.W., eds., *Tropical Deltas of Southeast Asia*: Sedimentology, Stratigraphy, and Petroleum Geology: Tulsa, Society of Sedimentary Geology (SEPM), p. 147-173.
- DASHTGARD, S.E., 2011a, Linking invertebrate burrow distributions (neoichnology) to physicochemical stresses on a sandy tidal flat: Implications for the rock record *Sedimentology*, v. 58, p. 1303-1325.
- DASHTGARD, S.E., 2011b, Neoichnology of the lower delta plain: Fraser River Delta, British Columbia, Canada: Implications for the ichnology of deltas: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 307, p. 98-108.
- GINGRAS, M.K., PEMBERTON, S.G., SAUNDERS, T., and CLIFTON, H.E., 1999, The ichnology of modern and Pleistocene brackish-water deposits at Willapa Bay, Washington: variability in estuarine settings: *Palaios*, v. 14, p. 352-374.
- HALES, W.J., 2000, The Impact of Human Activity on Deltaic Sedimentation, Marshes of the Fraser River Delta, British Columbia [unpublished Ph.D. thesis]: University of British Columbia, Vancouver, 178 p.
- HAM, D.G., 2005, Morphodynamics and sediment transport in a wandering gravel-bed channel: Fraser River, British Columbia [unpublished Ph.D. thesis]: University of British Columbia, Vancouver, 210 p.
- HEIN, F.J., and COTTERILL, D.K., 2006, The Athabasca Oil Sands - A regional geological perspective, Fort McMurray area, Alberta, Canada: *Natural Resources Research*, v. 15, p. 85-102.
- JOHNSON, S.M., 2012, The nature of inclined heterolithic stratification in a mixed tidal-fluvial setting: Depositional processes, sedimentology and ichnology, Middle Arm, Fraser River, Canada [unpublished M.Sc. thesis]: Simon Fraser University, Burnaby, 184 p.
- JOHNSON, W.A., 1921, Sedimentation of the Fraser River delta, *Geological Survey of Canada*, p. 46.
- KOSTASCHUK, R.A., and LUTERNAUER, J.L., 1989, The role of the salt-wedge in sediment resuspension and deposition: Fraser River estuary, Canada: *Journal of Coastal Research*, v. 5, p. 93-101.
- KOSTASCHUK, R.A., and ATWOOD, L.A., 1990, River discharge and tidal controls on salt-wedge position and implications for channel shoaling: Fraser River, British Columbia: *Canadian Journal of Civil Engineering*, v. 17, p. 452-459.
- KOSTASCHUK, R.A., and VILLARD, P., 1996, Flow and sediment transport overlarge subaqueous dunes: Fraser River, Canada: *Sedimentology*, v. 43, p. 849-863.
- KOSTASCHUK, R.A., and BEST, J., 2005, Response of sand dunes to variations in tidal flow: Fraser Estuary, Canada: *Journal of Geophysical Research*, v. 110, p. F04S04.
- KOSTASCHUK, R.A., LUTERNAUER, J.L., and CHURCH, M.A., 1998, Sedimentary processes in the estuary, *in* Clague, J.J., Luternauer, J.L., and Mosher, D.C., eds., *Geology and Natural Hazards of the Fraser River Delta*, British Columbia: Ottawa, Geological Survey of Canada, p. 41-56.
- KREISA, R.D., and MOIOLA, R.J., 1986, Sigmoidal tidal bundles and other tide-generated sedimentary structures of the Curtis Formation, Utah: *Geological Society of American Bulletin*, v. 97, p. 381-387.
- MACEACHERN, J.A., and GINGRAS, M.K., 2007, Recognition of brackish-water trace-fossil suites in the Cretaceous interior seaway of Alberta, Canada, *in* Bromley, R.G., Buatois, L.A., Mangano, G.M., Genise, J.F., and Melchor, R.N., eds., *Sediment-Organism Interactions: A Multifaceted Ichnology*: Tulsa, Oklahoma, SEPM, p. 149-193.
- MCLAREN, P., and REN, P., 1995, Sediment Transport and its environmental implications in the Lower Fraser River and Fraser delta, *in* Canada, E., ed., *Fraser River Action Plan*, p. 113.
- MCLEAN, D.G., 1990, The relation between channel stability and sediment transport on Lower Fraser River [unpublished Ph.D. thesis]: University of British Columbia, Vancouver, BC, 290 p.
- MCLEAN, D.G., CHURCH, M.A., and TASSONE, B., 1999, Sediment transport along the lower Fraser River. 1. Measurements and hydraulic computations: *Water Resources Research*, v. 35, p. 2533-2548.
- MILLIMAN, J.D., 1980, Sedimentation in the Fraser River and its estuary, southwestern British Columbia (Canada): *Estuarine and Coastal Marine Science*, v. 10, p. 609-633.
- MUSIAL, G., REYNAUD, J.-Y., GINGRAS, M.K., FENIES, H., LABOURDETTE, R., and PARIZE, O., 2012, Subsurface and outcrop characterization of large tidally influenced point bars of the Cretaceous McMurray Formation (Alberta, Canada) *Sedimentary Geology*, v. 279, p. 156-172.
- NELSON, A.D., and CHURCH, M., 2012, Placer mining along the Fraser River, British Columbia: The geomorphic impact: *GSA Bulletin*, v. 124, p. 1212-1228.
- PEARSON, N.J., and GINGRAS, M.K., 2006, An ichnological and sedimentological facies model for muddy point-bar deposits: *Journal of Sedimentary Research*, v. 76, p. 771-782.
- PEMBERTON, S.G., and WIGHTMAN, D.M., 1992, Ichnological characteristics of brackish water deposits, *in* Pemberton, S.G., ed., *Applications of Ichnology to Petroleum Exploration, A Core Workshop No.17*: Calgary, Alberta, SEPM, Society for Sedimentary Geology, p. 141-167.
- PUBLIC-WORKS-CANADA, 1957, The Fraser system, province of British Columbia: History of improvements 1887 to date.: New Westminster, Dominion of Public Works, p. 66.
- RANGER, M.J., and PEMBERTON, S.G., 1997, Elements of a stratigraphic framework for the McMurray Formation in south Athabasca area, Alberta, *in* Pemberton, S.G., and James, D.P., eds., *Petroleum Geology of the Cretaceous Mannville Group*, Western Canada: Calgary, Canadian Society of Petroleum Geologists, p. 263-291.
- SISULAK, C.F., and DASHTGARD, S.E., 2012, Seasonal controls on the development and character of inclined heterolithic stratification in a tide-influenced, fluvially dominated channel: Fraser River, Canada: *Journal of Sedimentary Research*, v. 82, p. 244-257.
- SMITH, D.G., 1988, Modern point bar deposits analogous to the Athabasca oil sands, *in* de Boer, P.L., van Gelder, A., and Nio, S.D., eds., *Tide-Influenced Sedimentary Environments and Facies*: Dordrecht, Springer, p. 417-432.
- SMITH, D.G., HUBBARD, S.M., LECKIE, D.A., and FUSTIC, M., 2009, Counter point bar deposits: lithofacies and reservoir significance in the meandering modern Peace River and ancient McMurray Formation, Alberta, Canada: *Sedimentology*, v. 56, p. 1655-1669.
- SMITH, D.G., HUBBARD, S.M., LAVIGNE, J.R., LECKIE, D.A., and FUSTIC, M., 2011, Stratigraphy of counter point bars and eddy accretion deposits in low energy meander belts of the Peace-Athabasca Delta, northeast Alberta, Canada, *in* Davidson, S.K., Leleu, S., and North, C.P., eds., *From River to Rock Record: The Preservation of Fluvial Sediments and Their Subsequent Interpretation*: Tulsa, OK, SEPM (Society for Sedimentary Geology), p. 143-152.
- STROBL, R.S., MUWAIS, W.K., WIGHTMAN, D.M., COTTERILL, D.K., and YUAN, L., 1997, Application of outcrop analogues and detailed reservoir characterization to the AOSTRA underground test facility, McMurray Formation, north eastern Alberta, *in* Pemberton, S.G., and James, D.P., eds., *Petroleum Geology of the Cretaceous Mannville Group*, Western Canada: Calgary, Canadian Society of Petroleum Geologists, p. 375-391.
- THOMSON, R.E., 1981, Oceanography of the British Columbia Coast: *Canadian Special Publication of Fisheries and Aquatic Sciences*, v. 56: Ottawa, Canada, Department of Fisheries and Oceans Canada, 291 p.
- VENDITTI, J.G., HUMPHRIES, R.P., ALLISON, M.A., NITTROUER, J.A., and CHURCH, M., 2010, Morphology and dynamics of a gravel-sand transition: 9th Federal Interagency Sedimentation Conference, p. 11.
- WCHL, 1977, Feasibility Study, Development of a Forty-Foot Draft Navigation Channel, New Westminster to Sandheads: Vancouver, Western Canada Hydraulic Laboratories, p. 107.

Accepted November 2012