

Supplementary Table 1. Extended list of publications (monographs/memoirs/article) with naticid gastropod listed in it. Qualitative site selectivity data were extracted from this list, because these papers did not quantify drilling gastropod predation. n = number of individuals (drilled + undrilled); DF = drilling frequency; IDF = incomplete DF; MULT = multiple DF; PE = prey effectiveness; – = no data provided. AP = Apertural, AB = abapertural, R = Random. AB^{\$}/AP^{\$} = site selectivity data extracted from figures but no quantifiable data provided in the original article, AB^m/AP^m = site selectivity mentioned in the article without providing raw data.

Time	Prey species	n	DF	Site	IDF (MULT)	PE	Size data	Country	References
Cretaceous	<i>Forator parkinsoni</i>	189	0.02	-	-	-	yes	UK	Taylor et al., 1983
	<i>Gyrodes gentii</i>	27	0.07	-	-	-	-	UK	Taylor et al., 1983
	Naticidae	304	0.02	-	0.01 (-)	0.33	-	USA	Kelley and Hansen, 2006
	<i>Euspira rectilabrum</i>	-	0.09	-	-	-	-	USA	Jones et al., 1998
	<i>Euspira rectilabrum</i>	-	0.01	-	-	-	-	USA	Jones et al., 1998
	<i>Euspira rectilabrum</i>	-	0.02	-	-	-	-	USA	Jones et al., 1998
	<i>Mammilla carnatica</i>	414	0.04	R	0.01 (0.00)	0.20	yes	India	Mallick et al., 2014
	<i>Mammilla</i> sp.	2	0.00	-	0.00 (0.00)	0.00	yes	India	Mallick et al., 2014
	<i>Euspira obliquata</i>	403	0.02	AP ^{\$}	-	-	-	USA	Li et al., 2011
Paleocene	Naticidae	18	0.17	-	0.00(-)	0.00	-	USA	Kelley and Hansen, 2006
	<i>Polinices clementensis</i>	-	-	AP ^{\$}	- (-)	-	-	USA	Hickman, 1980
	Naticidae	61	0.10	-	0.00(-)	0.00	-	USA	Kelley and Hansen, 2006
Eocene	<i>Natica epiglottina</i>	31	0.16	-	0.00 (-)	0.00	-	France	Taylor, 1970
	<i>Sinum clathratus</i>	06	0.17	-	0.00 (-)	0.00	-	France	Taylor, 1970

	<i>Natica obscura</i>	65	0.09	AP	0.00 (0.00)	0.00	-	India	JUM collection
	<i>Natica amekiensis</i>	238	0.14	AB ^m	0.05 (-)	0.27	yes	Nigeria	Adegoke and Tevesz, 1974
	<i>Polinices marambioensis</i>	2648	0.11	AP	0.04 (0.04)	0.27	yes	Antarctica	Brezina et al., 2016
	Naticidae	74	0.08	-	0.00 (-)	0.00	-	USA	Kelley and Hansen, 2006
	Naticidae	228	0.14	-	0.04 (-)	0.22	-	USA	Kelley and Hansen, 2006
	Naticidae	54	0.15	-	0.02 (-)	0.11	-	USA	Kelley and Hansen, 2006
	<i>Carinacca rosensis</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Euspira nuciformis</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Polinices clementensis</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Tejonia meragai</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Euspira labellata</i>	115	0.08	AP ^m	0.09 (-)	0.53	-	France	Taylor, 1970
	<i>Natica</i> spp.	395	0.21	-	0.00 (0.00)	0.00	-	France	Fischer, 1966
Oligocene	<i>Natica helicina</i>	140	0.49	AB ^m	-	-	-	Germany	Klähn, 1932
	Naticidae	342	0.10	-	0.00 (-)	0.00	-	USA	Kelley and Hansen, 2006
	Naticidae	40	0.15	-	0.00 (-)	0.00	-	USA	Kelley and Hansen, 2006
	<i>Polinices helicina</i>	-	-	-	- (0.10)	-	-	Hungary	David, 1999

	<i>Polinices josephina</i>	-	-	-	- (0.33)	-	-	Hungary	David, 1999
	<i>Natica tigrina</i>	-	-	-	- (0.03)	-	-	Hungary	David, 1999
	<i>Natica helicina</i>	209	0.33	AB ^m	- (-)	-	-	Germany	Klähn, 1932
	Naticidae	-	-	-	-	-	-	Hungary	Balaska, 2014
Miocene	Naticidae	-	-	AP ^s	-	-	-	Japan	Majima, 1989
	Naticidae	-	-	AP ^s	-	-	-	USA	Addicott, 1970
	<i>Naticarius posuncula</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Naticarius teglandea</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Euspira lincolnensis</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Cryptonatica oregonensis</i>	150	0.03	-	-	-	-	USA	Colbath, 1985
	<i>Cryptonatica oregonensis</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	<i>Tectonatica janthostoma</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	Naticidae	1402	0.29	-	0.01 (-)	0.03	-	USA	Kelley and Hansen, 2006
	<i>Lunatia heros</i>	530	0.17	AP	0.03 (-)	0.17	Yes	USA	Kelley, 1991
	<i>Polinices duplicatus</i>	340	0.09	R	0.02 (-)	0.19	yes	USA	Kelley, 1991
	<i>Natica helloina</i>	189	0.09	-	-	-	-	Poland	Hoffman et al., 1974
	<i>Natica joshephina</i>	202	0.11	-	-	-	-	Poland	Hoffman et al., 1974

	<i>Natica millepunctata</i>	428	0.11	-	-	-	-	Poland	Hoffman et al., 1974
	<i>Natica redempta</i>	159	0.09	-	-	-	-	Poland	Hoffman et al., 1974
	<i>Natica pseudoredempta</i>	81	0.18	-	-	-	-	Poland	Hoffman et al., 1974
	<i>Natica</i> spp.	196	0.09	-	-	-	yes	Poland	Kowalewski, 1990
	<i>Natica</i> spp.	14	0.65	-	-	-	-	Panama	Fortunato, 2007
	<i>Polinices</i> spp.	26	0.25	-	-	-	-	Panama	Fortunato, 2007
	<i>Stigmaulux</i> spp.	6	1.00	-	-(-)	-	-	Panama	Fortunato, 2007
	<i>Natica obscura</i>	296	0.04	R	0.00 (0.00)	0.00	yes	India	Goswami et al., 2015
	Naticidae	403	0.13	—	0.01 (-)	0.07	-	Austria, Slovenia	Sawyer and Zuschin, 2011
	<i>Neverita duplicata</i>	1472	0.11	-	0.01 (-)	0.08	yes	USA	Dietl and Alexander, 1995, 2000
	<i>Euspira heros</i>	1163	0.17	-	0.01 (-)	0.05	yes	USA	Dietl and Alexander, 1995, 2000
	<i>Natica tigrina</i>	1826	0.05	-	0.01 (-)	0.17	-	Poland	Zlotnik, 2001
	<i>Taniella mima</i>	123	0.19	-	-	-	-	New Zealand	Maxwell, 1988
	<i>Friginatica vaughani</i>	40	0.03	-	-	-	-	New Zealand	Maxwell, 1988
	<i>Polinices redempta</i>	14	0.07	-	- (-)	-	-	Slovakia	Pek et al., 1997
	<i>Nacca unica</i>	2	-	AP ^s	- (-)	-	-	Poland	Baluk, 2006
	<i>Euspira</i> sp.	97	0.16	-	0.00 (0.00)	0.00	-	Europe	Hoffmeister, 2002

	<i>Natica</i> sp.	21	0.05	-	0.00 (0.00)	0.00	-	Europe	Hoffmeister, 2002
Pliocene	Naticidae	-	-	AP ^{\$}	-	-	-	Japan	Majima, 1989
	<i>Natica scethra</i>	39	~0.5 2	-	-	-	yes	Ecuador	Walker, 2001
	<i>Natica millepunctata</i>	-	-	R ^{\$}	-	-	-	Italy	Bromley, 1981
	<i>Natica tigrina</i>	32	0.34	-	-	-	-	Spain	Hoffman and Martinell, 1984
	<i>Tectonatica tectula</i>	30	0.20	-	-	-	-	Spain	Hoffman and Martinell, 1984
	<i>Lunatia helicina</i>	94	0.19	-	-	-	-	Spain	Hoffman and Martinell, 1984
	Naticidae	-	-	AB ^{\$}	-	-	-	Netherlands	Klompmaker, 2009
	Naticidae	448	0.20	-	0.00 (-)	0.00	-	USA	Kelley and Hansen, 2006
	<i>Polinices</i> sp.	191	0.07	-	-	-	-	Belgium	Boekschoten, 1967
	<i>Polinices cumminghamensis</i>	205	0.01	-	-	-	-	Australia	Hingston, 1985
	<i>Cochlis pseudoepiglottina</i>	183	0.45	AP ^{\$}	0.13 (-)	0.22	yes	France	Martinell et al., 2010
	<i>Naticarius canrena</i>	65	0.12	R	0.00	0.00	yes	USA	Christie, 2009
	<i>Neverita duplicata</i>	170	0.05	R	0.00 (0.00)	0.00	yes	USA	Christie, 2009

	<i>Euspira heros</i>	159	0.11	R	0.00 (0.00)	0.00	yes	USA	Christie, 2009
	<i>Lunatia perspectiva</i>	41	0.10	R	0.00 (0.00)	0.00	yes	USA	Christie, 2009
	<i>Tectonatica pusilla</i>	28	0.14	R	0.00 (0.00)	0.00	yes	USA	Christie, 2009
	<i>Lunatia interna</i>	94	0.22	R	0.00 (0.00)	0.00	yes	USA	Christie, 2009
	<i>Polinices</i> sp.	11	0.27	AP	0.00 (0.00)	0.00	yes	USA	Christie, 2009
Pleistocene	<i>Natica chemnitzii</i>	-	~0.2 0	AB ^m	-	-	-	Mexico	Stump, 1975
	<i>Neverita reclusiana</i>	-	-	AP ^s	- (-)	-	-	USA	Marincovich, 1977
	Naticidae	-	-	AP ^s	-	-	-	Japan	Majima, 1989
	<i>Natica arachnoidea</i>	33	0.33	R	-	-	yes	Fiji	Kabat and Kohn, 1986
	<i>Eunaticina linnaeana</i>	06	0.17	-	-	-	yes	Fiji	Kabat and Kohn, 1986
	<i>Natica gualteriana</i>	34	0.30	R	-	-	yes	Fiji	Kabat and Kohn, 1986
	<i>Natica areolata</i>	21	0.38	R	-	-	yes	Fiji	Kabat and Kohn, 1986
	<i>Natica euzona</i>	47	0.30	R	-	-	yes	Fiji	Kabat and Kohn, 1986
	<i>Polinices peselephantii</i>	29	0.03	-	-	-	yes	Fiji	Kabat and Kohn, 1986

	<i>Natica</i> sp.	07	0.00	-	-(-)	-	yes	Fiji	Kabat and Kohn, 1986
	Naticidae	312	0.19	-	~0.00 (-)	0.00	-	USA	Kelley and Hansen, 2006
	<i>Neverita duplicata</i>	501	0.17	-	0.01 (-)	0.06	-	USA	Dietl and Alexander, 1995, 2000
	Naticidae	207	0.19	-	-	-	yes	Fiji	Kohn and Arua, 1999
	<i>Neverita duplicata</i>	755	0.14	R	~0.00 (~ 0.00)	0.00	-	USA	Christie, 2009
	<i>Euspira heros</i>	327	0.16	R	0.00 (0.00)	0.00	-	USA	Christie, 2009
	<i>Naticarius canrena</i>	126	0.06	AP	0.00 (0.00)	0.00	-	USA	Christie, 2009
	<i>Tectonatica pusilla</i>	153	0.05	R	0.00 (0.00)	0.00	-	USA	Christie, 2009
Holocene	<i>Natica gualteriana</i>	1773	0.55	AP	~0.00 (0.00)	0.00	yes	Indian Subcontinents	Das et al., 2014
	<i>Natica tigrina</i>	857	0.57	AP	0.01 (0.02)	0.01	yes	India	Pahari et al., 2016
	<i>Tectonatica janthostoma</i>	-	-	AP ^{\$}	- (-)	-	-	Japan	Marincovich, 1977
	<i>Natica lineata</i>	760	0.01	R	~0.00 (0.00)	0.00	yes	India	JUM collection

	<i>Polinices mammilla</i>	55	0.00	-	0.00 (0.00)	0.00	yes	India	JUM collection
	<i>Polinices didyma</i>	619	0.06	AP	~0.00 (0.00)	0.00	yes	India	JUM collection
	<i>Eunaticina</i> sp.	296	0.13	AP	0.00 (0.00)	0.00	yes	India	JUM collection
	<i>Sinum neritoidium</i>	15	0.00	-	0.00 (0.00)	0.00	yes	India	JUM collection
	<i>Polinices</i> sp.	08	0.13	-	-(-)	-	-	Bahamas	Pruss et al., 2011
	Naticidae	10	0.41	-	- (-)	-	-	USA	Funderburk, J. 2010
	<i>Euspira heros</i>	-	-	R	- (-)	-	-	Canada	Clements, 2011
	<i>Polinices sordidus</i>	220	0.02	-	-	-	-	Australia	Green, 1968
	<i>Polinices conicus</i>	112	0.15	-	-	-	-	Australia	Green, 1968
	<i>Euspira heros</i>	225	0.13	R	0.10 (-)	0.43	-	USA	Dietl and Alexander, 1995, 2000
	<i>Neverita duplicata</i>	736	0.10	R	0.32 (-)	0.24	-	USA	Dietl and Alexander, 1995, 2000

REFERENCES

- ADEGOKE, O.S., and TEVESZ, M.J.S., 1974, Gastropod predation patterns in the Eocene of Nigeria: *Lethaia*, v. 7, p. 17-24.
- ADDICOTT, W.O., 1970, Miocene gastropods and biostratigraphy of the Kern River area, California: Geological survey professional paper 642.
- ALEXANDER, R.R., and DIETL, G.P., 2001, Shell repair frequencies in New Jersey bivalves: a recent baseline for tests of escalation with Tertiary, Mid-Atlantic congeners: *Palaios*, v. 16, p. 354-371.
- ALLMON, W.D., NIEH, J.C., and NORRIS, R.D., 1990, Drilling and peeling of turritelline gastropods since the Late Cretaceous: *Palaeontology*, v. 33, p. 595-611.
- BALASKA, P., 2014. Gastropod Predation on Late-Oligocene Age Molluscs – a Comparison (Molluscan Clay, Wind Brickyard, Eger). 8th International Bioerosion Workshop, Eger, Hungary.
- BALUK, W., 2006, Middle Miocene (Badenian) gastropods from Korytnica, Poland; Part V, Addenda et Corrigenda ad Prosobranchia: *Acta Geologica Polonica*, v. 56, p. 177-220.
- BANDEL, K., 1999. On the origin of the carnivorous gastropod groups Naticoidea (Mollusca) in the Cretaceous with description of some convergent but unrelated groups: *greifswalder Geowissenschaftliche Beiträge*, v. 6, p. 143-175.
- BARDHAN, S., DAS, S.S., MALLICK, S., MONDAL, S., and DUTTA, R., 2012.a, The record of oldest naticid (gastropod from the Oxfordian (Upper Jurassic) of Kutch, India: palaeoecological implications: Geological Society of America Abstract with program, v. 44(7).

- BARDHAN, S., CHATTOPADHYAY, D., MONDAL, S., DAS, S. S., MALLICK, S., ROY, A., and CHANDA, P., 2012.b, Record of intense predatory drilling from Upper Jurassic bivalves of Kutch, India: Implications for the history of biotic interaction: Palaeogeography Palaeoclimatology Palaeoecology, v. 317–318, p. 153-161.
- BROWN, G.M., MONDAL, S., HARRIES, P.J., and SLATTERY, J.S., 2014, Drilling predation patterns from the Miocene Chipola formation of Florida: taphonomic versus environmental overprinting: Geological Society of America Abstract with program, v. 46(6).
- BOEKSHOTEN, G.J., 1967, Palaeoecology of some Mollusca from the Tielrode Sands (Pliocene, Belgium): Palaeogeography Palaeoclimatology Palaeoecology, v. 3, p. 311-362.
- CHATTOPADHYAY, D., SARKAR, D., and DUTTA, S., 2014, What controls cannibalism in drilling gastropods? A case study on *Natica tigrina*: Palaeogeography Palaeoclimatology Palaeoecology, v. 410, p.126-133.
- CHRISTIE, M., 2009. Ecological interactions across a Plio-Pleistocene interval of faunal turnover: Naticid cannibalism north and south of Cape Hatteras, North Carolina: PhD Dissertation, The College of William and Mary, USA.
- CLEMENTS, J.C., 2011. Feeding Ecology of the Northern Moonsnail (*Euspira heros*)(Gastropoda: Naticidae) in Cape Breton, Nova Scotia: Dissertation, Cape Breton University.
- COLBATH, S.L., 1985, Gastropod predation and depositional environment of two molluscan communities from the Miocene Astoria Formation at Beverly Beach State Park, Oregon: Journal of Paleontology, v. 59, p.849–869.
- DAS, A., MONDAL, S., and BARDHAN, S., 2014, A note on exceptionally high confamilial naticid drilling frequency on *Natica gualteriana* from the Indian subcontinent: Historical Biology, v. 26, p.758-764.

DAVID, A., 1999, Predation by Naticid Gastropods on Late-Oligocene (Egerian) Molluses Collected from Wind Brickyard, Eger, Hungary: Malakologiatajekoz tat o Malacological Newsletter, v. 17, p.11–19.

DIETL, G.P., and ALEXANDER, R.R., 1995, Borehole site and prey size specificity in naticid predation on *Euspira (Lunatia) heros* Say and *Neverita (Polinices) duplicata* Say from the southern New Jersey Coast: Journal of Shellfish Research, v. 14, p. 307-314.

DIETL, G.P., and ALEXANDER R.R., 2000, Post-Miocene shift in stereotypic naticid predation on confamilial prey from the mid-Atlantic shelf: Coevolution with dangerous prey: *Palaios*, v. 15, p. 414–429.

DIETL, G.P., KELLEY, P.H., 2001, Mid-Paleozoic latitudinal predation gradient: Distribution of brachiopod ornamentation reflects shifting Carboniferous climate: *Geology*, v. 29, p. 111-114.

DUDLEY, E.C., and VERMEIJ, G.J., 1978, Predation in time and space: drilling in the gastropod *Turritella*: *Paleobiology*, v. 4, p. 436-441.

FISCHER, P.H., 1966, Perforations de fossiles Tertiaires par des gastéropodes prédateurs. *J. Conchol.*, v. 105, p. 66-96.

FUNDERBURK, J., 2010, Modern Variation in Predation Intensity: Constraints on Assessing Predator-Prey Relationships in Paleoecologic Reconstructions: PhD Thesis, University of South Florida, Tampa, USA.

FUTUYMA, D.J., and SLATKIN, M. 1983, Coevolution: Sinauer Associates Inc., Sunderland, 555 p.

GOSWAMI, P., DAS, S. BARDHAN,S., MALLICK, S., S. and PAUL, S., 2015, Drilling gastropod predation on the lower Miocene gastropod assemblages from Kutch, western India: spatiotemporal implications: *Geological Society of America Abstract with Program*, v. 47, p. 426.

GREEN, R.H., 1968. Mortality and stability in a low diversity subtropical intertidal community: *Ecology*, v. 49, p. 848-854.

HANSEN, T. A., and KELLEY, P.H., 1995, Spatial variation of naticid gastropod predation in the Eocene of North America: *Palaios*, v. 10: p. 268-278.

- HANSEN, T.A., KELLEY, P.H., MELLAND, V.D., and GRAHAM, S.E., 1999, Effect of climate-related mass extinctions on escalation in molluscs: *Geology*, v. 27, p. 1139-1142.
- HARRIES, P.J., and SCHOPF, K.M., 2007, Late Cretaceous gastropod drilling intensities: data from the Maastrichtian Fox Hills Formation, western interior seaway, USA: *Palaios*, v. 22, p. 35-46.
- HIGSTON, J. P., 1985, Predation patterns among molluscs in the Victorian Tertiary. *Proceedings of the Royal Society of Victoria*, v. 97, p. 49-57.
- HOFFMAN, A. PISERA, A., and RYSZKIEWICZ, M., 1974, Predation by muricid and naticid gastropods on the lower Tortonian mollusks from the Korytnica clays: *Acta Geologica Polonica*, v. 24, p. 249-264.
- HOFFMAN, A., and MARTINELL, J., 1984, Prey selection by naticid gastropods in the Pliocene of Emporda (Northeast Spain): *NeusJahrbuch fur Geologie und Paaontologie*, v. 7, p. 393-399.
- HOFFMEISTER, A.P., 2002, Quantitative analysis of drilling predation patterns in the fossil record: ecological and evolutionary implications: PhD Dissertation, Virginia Polytechnic Institute and State University, USA.
- HELENA-FORTUNATO, H., 2007, Naticid gastropod predation in the Gatun Formation (late middle Miocene), Panama: Preliminary assessment: *Palaeontologische Zeitschrift*, v. 81, p. 356–364.
- HUNTLEY, J.W., and KOWALEWSKI, M., 2007, Strong coupling of predation intensity and diversity in the Phanerozoic fossil record: *Proceedings of the National Academy of Sciences*, v. 104, p. 15006-15010.
- HUTCHINGS, J.A., and HERBERT, G.S., 2013, No honor among snails: conspecific competition leads to incomplete drill holes by a naticid gastropod: *Palaeogeography Palaeoclimatology Palaeoecology*, v. 379, p. 32-38.

- JONES, M.A., KELLEY, P.H., ERICKSON, J.M., and HANSEN, T.A., 1998, Interregional comparison of drilling on Cretaceous naticid gastropods and implications for the hypothesis of escalation: Geological Society of America Abstracts with Programs, v. 30, p. 25.
- KABAT, A.R., and KOHN, A.J., 1986. Predation on Early Pleistocene naticid gastropods in Fiji: Palaeogeography Palaeoclimatology Palaeoecology, v. 53, p. 255–269.
- KELLEY, P. H., 1991. Apparent cannibalism by Chesapeake Group naticid gastropods: A predictable result of selective predation: Journal of Paleontology, v. 65, p. 75-79.
- KELLEY, P.H., HANSEN, T.A., GRAHAM, S.E., and HUNTOON, A.G., 2001, Temporal patterns in the efficiency of naticid gastropod predators during the Cretaceous and Cenozoic of the United States Coastal Plain: Palaeogeography Palaeoclimatology Palaeoecology, v. 166, p. 165-176.
- KELLEY, P. H., and HANSEN, T. A., 2003, The fossil record of drilling predation on bivalves and gastropods. In: Kelley PH, Kowalewski M, Hansen TA (eds) Predator-Prey Interactions in the Fossil Record: Topics in Geobiology Series 20, Plenum Press/Kluwer, New York.
- KELLEY, P.H., and HANSEN, T.A., 1996, Naticid gastropod prey selectivity through time and the hypothesis of escalation: Palaios, v. 11, p. 437-445.
- KELLEY, P.H., and HANSEN, T.A., 2006, Comparisons of class- and lower taxon-level patterns in naticid gastropod predation, Cretaceous to Pleistocene of the US Coastal Plain: Palaeogeography Palaeoclimatology Palaeoecology, v. 236, p. 302–320.
- KELLEY, P.H., and HANSEN, T.A., 2007, A case for cannibalism: confamilial and conspecific predation by naticid gastropods, Cretaceous through Pleistocene of the United States Coastal Plain, in KELLEY, P.H., and HANSEN, T.A., eds., Predation in Organisms: Springer Berlin Heidelberg, p. 151-170.

KITCHELL, J.A., BOGGS, C.H., KITCHELL, J.F., and RICE, J.A., 1981, Prey selection by naticid gastropods: experimental tests and application to application to the fossil record: *Paleobiology*, v. 7, p. 533-552.

KLÄHN, H.A.N.S., 1932, Der Bohrakt fossiler bohrender Schnecken und das Vernichtungsmass durch räuberische Gastropoden des Sternberger Oligozäns: *Sitzungsber. u. Abh. der Naturf. Gesellschaft zu Rostock*, v. 3, p. 89-103.

KOHN, A.J., and ARUA, I., 1999, An Early Pleistocene molluscan assemblage from Fiji: Gastropod faunal composition, paleoecology and biogeography: *Palaeogeography Palaeoclimatology Palaeoecology*, v. 146, p. 99–145.

KOWALEWSKI, M., 1990, A hermeneutic analysis of the shell-drilling gastropod predation on mollusks in the Korytnica Clays (Middle Miocene; Holy Cross Mountains, Central Poland): *Acta Geologica Polonica*, v. 40, p. 183-213.

KOWALEWSKI, M., DULAI, A., and FÜRSICH, F.T., 1998, A fossil record full of holes: the Phanerozoic history of drilling predation: *Geology*, v. 26, p. 1091-1094.

KLONPMAKER, A., 2009, Taphonomic bias on drill-hole predation intensities and paleoecology of Pliocene mollusks from Langenboom (Mill), The Netherlands: *Palaios*, v. 24, p. 772–779.

LI, R.Y., YOUNG, H.R., and ZHAN, R.B., 2011, Drilling predation on scaphopods and other molluscs from the Upper Cretaceous of Manitoba, Canada: *Palaeoworld*, v. 20, p.296–307.

MAJIMA, R., 1989, Cenozoic fossil Naticidae (Mollusca: Gastropoda) in Japan.

MAXWELL, P.A., 1988. Late Miocene deep-water mollusca from the Stillwater Mudstone at Greymouth, Westland, New Zealand: New Zealand Geological Survey Paleontological Bulletin, 120 pp.

- MARTINELL, J., DOMENECH, R., AYMAR, J., and KOWALEWSKI, M., 2010, Confamilial predation in Pliocene naticid gastropods from southern France: utility of preexisting collections in quantitative paleoecology: *Palaios*, v. 25, p. 221-228.
- MONDAL, S., HERBERT, G., 2012, Effect of extinction on durophagy: a case study from the Neogene of Florida: Geological Society of America Abstract with program, v. 44(7).
- MONDAL, S., and HARRIES, P., 2013, Prey selectivity by the predatory drilling gastropods: a Meso-Cenozoic perspective: Geological Society of America Abstract with program, v. 45(7), p. 315.
- MONDAL, S., and HARRIES, P.J., 2015, Temporal Patterns in Successful and Unsuccessful Shell-Breaking Predatory Attack Strategies on *Varicorbula* in the Plio-Pleistocene of Florida: *Palaeogeography Palaeoclimatology Palaeoecology*, v. 428, p. 31-38.
- MONDAL, S., HUTCHINGS, J.A., and HERBERT, G.S., 2014, A Note on Edge Drilling Predation by Naticid Gastropods: *Journal of Molluscan Studies*, v. 80, p. 206–212.
- PAHARI, 2015, Subaerial drilling predation by *Natica tigrina* on the intertidal molluscan community of Chandipur, eastern coast of India: Jadavpur University MS Thesis, Unpublished.
- PAUL, G., DAS. A., BARDHAN, S., and MONDAL, S., 2013, Predation on Recent turritelline gastropods from the Indian subcontinent and comparison with a revised global database: *Malacologia*, v. 56, p. 193-213.
- PEK, I., MIKULÁŠ, R., LYSÁKOVÁ, G., 1997, Boring ichnofossils on mollusc shells from the Late Badenian at Rohožník (Malé Karpaty Mts., Slovakia): *Zemní Plyn a Nafta*, v. 42, p. 47-55.
- PRUSS, S. B., STEVENSON, M., and DUFFEY, S., 2011, Drilling predation and taphonomy in modern mollusk death assemblages, San Salvador Island, Bahamas: *Palaeogeography Palaeoclimatology Palaeoecology*, v. 311, p. 74–81.

SARKAR, D., BARDHAN, S., MONDAL, S., DAS, A., PAHARI, A., BURAGOHAIN, D., and SAHA, S., in press. Predation on Recent terebrid gastropods from the Indian Subcontinents and a spatiotemporal reappraisal based on a revised global database: *Malacologia*.

SAWYER, J. A., and ZUSCHIN, M., 2011, Drilling predation in mollusks from the Lower and Middle Miocene of the Central Paratethys: *Palaios*, v.26, p. 284–297.

SMITH, A.G., SMITH, D. G., and FUNNELL, B. M., 1994, *Atlas of Mesozoic and Cenozoic coastlines*. Cambridge University Press, Great Britain.

SOHL, N.F., 1969. The fossil record of shell boring by snails: *American Zoologist*, v. 9, p. 725-734.

STUMP, T.E., 1975, Pleistocene molluscan paleoecology and community structure of the Puerto Libertad region, Sonora, Mexico: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 17, p. 177-226.

TAYLOR, J. D., 1970, Feeding habits of predatory gastropods in a Tertiary (Eocene) molluscan assemblage from the Paris basin: *Palaeontology*, v. 13, p. 255-260.

TAYLOR, J.D., CLEEVELY, R.J., and MORRIS, N.J., 1983, Predatory gastropods and their activities in the Blackdown Greensand (Albian) of England: *Palaeontology*, v. 26, p. 521–553.

VERMEIJ, G.J., 1987, *Evolution and escalation: an ecological history of life*. Princeton University Press. Princeton, 544 p.

VISAGGI, C.C., and KELLEY, P.H., 2015, Equatorward increase in naticid gastropod drilling predation on infaunal bivalves from Brazil with paleontological implications: *Palaeogeography Palaeoclimatology Palaeoecology*, v. 438, p. 285-299.

WALKER, S.E., 2001, Paleoecology of gastropods preserved in turbiditic slope deposits from the upper Pliocene of Ecuador: *Palaeogeography Palaeoclimatology Palaeoecology*, v. 166, p. 141–163.

WHORLEY, T.L., MONDAL, S., HARRIES, P.J., and SLATTERY, J.S., 2014, Impact of environmental variability on drilling predation: a comparison among *Turritella*-dominated assemblages: Geological Society of America Abstract with program, v. 46(6).

ZLOTNIK, M., 2001. Size-related changes in predatory behaviour of naticid gastropods from the Middle Miocene Korytnica Clays, Poland: Acta Palaeontologica Polonica, v. 46, p. 87-97.

Supplementary Table 2. List of additional references scanned to extract additional data on predation intensities and site selectivity of naticid confamilial predation. These resources listed naticid gastropod but did not mention predation intensity or site selectivity of confamilial naticid predation data.

1. Allison, E.C. 1955. Middle Cretaceous Gastropoda from Punta China, Baja California, Mexico. *Journal of Paleontology*. 1: 400-432.
2. Ansell, A.D., Macé, A.M. 1978. Comparative studies of the gastropod, *Polinices alderi* from Mediterranean and North Atlantic populations. *Haliotis*. 9: 65-72.
3. Bandel, K. 1999. On the origin of the carnivorous gastropod group Naticoidea (Mollusca) in the Cretaceous with description of some convergent but unrelated groups. *Greifswalder Geowissenschaftliche Beiträge*. 6: 143-175.
4. Beistro, F.I., Medcof, J., Lord, I. 1965. Clam drill (*Polinices*) investigations at St Andrews. 1948. *Manuscript Reports of the Biological Stations, Report 845*.
5. Berg, C.J. 1976. Ontogeny of Predatory Behavior in Marine Snails (Prosobranchia: Naticidae). *The Nautilus*. 90: 1-4.
6. Bryan and Jones.1989. Fabric of the Cretaceous -Tertiary marine macrofaunal transition at Braggs, Alabama. *Paleo3*
7. Chattopadhyay, D. 2009. Predation in molluscs: A multi-taxon approach using neontological and paleontological data. ProQuest.
8. Chattopadhyay, D., Sarkar, D., Dutta, S., Prasanjit, S.R. 2014. What controls cannibalism in drilling gastropods? A case study on *Natica tigrina*. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 410: 126-133.
9. Eames, F.E. 1952. A contribution to the study of the Eocene in western Pakistan and western India C. The description of the Scaphopoda and Gastropoda from Standard sections in the Rakhi Nala and Zinda Pir areas of the western Punjab and in the Kohat district. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*. 107:1-168.
10. Gardner, 1926. The molluscan fauna of the Alum Bluff Group of Florida. No. 142. US Government Printing Office.
11. Gardner, J.A., Vaughan, T.W., Popenoe, W.P. 1935. The Midway Group of Texas. The University.
12. Gardner, J.A., Mansfield, W.C. 1943. Mollusca from the Miocene and lower Pliocene of Virginia and North Carolina. No. 199. US Government Printing Office.
13. Gould, E.S. 2010. Unexpected rates of cannibalism under competitive conditions by the naticid gastropod *Neverita duplicata* (Say). Diss. U. North Carolina Wilmington.
14. Griffin, M., Hünicken, M.A. 1994. Late Cretaceous-Early Tertiary gastropods from southwestern Patagonia, Argentina. *Journal of Paleontology*. 68: 257-274.
15. Harzhauser, M., 2014. A seagrass-associated Early Miocene Indo-Pacific gastropod fauna from South-West India (Kerala). *Palaeontographica Abteilung. A*: 73-178.
16. Harzhauser, M., Reuter, M., Piller, W.E., Berning, B., Kroh, A., Mandic, O. 2009. Oligocene and Early Miocene gastropods from Kutch (NW India) document an early biogeographic switch from Western Tethys to Indo- Pacific. *Paläontologische Zeitschrift*. 83: 333-372.
17. Heidelberger, D., Amher, M.R. 2002. Devonian Gastropoda from the Dornap "Massenkalk" complex (Bergisches Land, Germany). *Paläontologische Zeitschrift*. 76: 317-329.
18. Hislop, S., 1860. On the Tertiary deposits, associated with trap-rock, in the East Indies with descriptions of the Fossil shells, by the Rev. S. Hislop; and of the fossil insects, by Andrew Murray, Esq., FRSE; and a note on the fossil Cypridæ, by T. Rupert Jones, Esq., FGS. *Quarterly Journal of the Geological Society*, 16: 154-166.
19. Hoffman, A., 1978. Character shift in the naticid gastropods from the Badenian (Miocene) of Poland. *Acta Palaeontologica Polonica*, 23: 31-39.

20. Jain, R.L. 2014. Neogene fossils from Kathiawar, Gujarat, India with special emphasis on taxonomic description of molluscs and corals. Geological Survey of India.
21. Jaitly, A.K., Fürsich, F.T., Heinze, M. 1995. Contributions to the Jurassic of Kachchh, western India. IV. The bivalve fauna. Part I. Subclasses Palaeotaxodonta, Pteriomorpha, and Isofilibranchia. *Beringeria*. 16: 147-257.
22. Kabat, A.R. 2000. Results of the Rumphius Biohistorical Expedition to Ambon (1990). Part 10. Mollusca, Gastropoda, Naticidae. *Zoologische Mededelingen*. 73: 345-380.
23. Kaim, A., Jenkins, R.G., Hikida, Y. 2009. Gastropods from Late Cretaceous Omagari and Yasukawa hydrocarbon seep deposits in the Nakagawa area, Hokkaido, Japan. *Acta Palaeontologica Polonica*. 54: 463-490.
24. Kase, T. 1984. Early Cretaceous marine and brackish-water Gastropoda from Japan. *National Science Museum Monographs*. 1: 1-189.
25. Kensley, B.F., Pether, J. 1986. Late tertiary and early quaternary fossil mollusca of the Hondeklip area, Cape Province, South Africa. 97. South African Museum.
26. Khosla, S.C. 2000. Atlas of Early Palaeogene invertebrate fossils of the Himalayan foothills belt. *Geological Society of India*. 56: 229-230.
27. Kulkarni, K.G., Kapoor, S.B., Borkar, V.D. 2010. Molluscan fauna from the Miocene sediments of Kachchh, Gujarat, India—Part 3. Gastropods. *Journal of earth system science*. 119: 307-341.
28. MacNeil, F.S., 1964. Eocene megafossils from Ishigaki-shima, Ryukyu-retto. No. 399-B.
29. Mansfield, W.C. 1930. Miocene gastropods and scaphopods of the Choctawhatchee Formation of Florida. *Bulletin 3. Florida state Geological Survey*.
30. Marwick, J. 1965. Upper Cenozoic Mollusca of Wairoa District, Hawke's Bay. *New Zealand Department of Scientific and Industrial Research, New Zealand Geological Survey*.
31. Marwick, J. 1931. The tertiary mollusca on the Gisborne district. *Paleontological Bulletin*. NZ Geological survey. 13:1-177.
32. Petuch,E.J. *Atlas of Florida Fossil Shells.(Pliocene and Pleistocene Marine Gastropods)*.
33. Popenoe, W.P., Saul, L.R., Susuki, T. 1987. Gyroform gastropods from the Pacific coast Cretaceous and Paleocene. *Journal of Paleontology*. 61: 70-100.
34. delRio, C.J. 2014. Tertiary Marine Molluscan Assemblages of Eastern Patagonia (Argentina): A Biostratigraphic Analysis. *Journal of Paleontology*. 78: 1097-1122.
35. del Rio, C.J. 2012. A New Early Danian Gastropod Assemblage from Northern Patagonia, Río Negro Province, Argentina. *Journal of Paleontology*. 86:1002-1016
36. Savazzi, E., Reymert, R.A. 1989. Subaerial hunting behaviour in *Natica gualteriana* (naticid gastropod). *Palaeogeography, palaeoclimatology, palaeoecology*. 74: 355-364.
37. Schnetler, K. I. 2001. The Selandian (Paleocene) mollusc fauna from Copenhagen, Denmark: the Poul Harder 1920 collection. *Geology of Denmark Survey Bulletin*. No. 3 7.
38. Sohl, N.F. 1965. Marine Jurassic gastropods, central and southern Utah. No. 503-D. US Government Printing Office.
39. Sohl, N.F. 1967. Upper Cretaceous gastropods from the Pierre Shale at Red Bird, Wyoming. US Government Printing Office.
40. Stoliczka, F. 1867. Cretaceous fauna of Southern India II — the Gastropoda. *Memoirs of the geological survey of India. Palaeontologia Indica*. No. 2.
41. Thomas A.D., Kendrick, G.W. 2008. Silicified Eocene molluscs from the Lower Murchison district, Southern Carnarvon Basin, Western Australia. *Records of the Western Australian Museum*. 24: 217-246.
42. Vredenburg, E.W. 1928. Descriptions of mollusca from the Post-Eocene Tertiary formation of North-Western India: Gastropoda (in part) and lamellibranchiate. *Memoir of Geological Survey of India*. 50: 351-506.
43. Wade, B. 1926. The fauna of the Ripley Formation on Coon Creek, Tennessee. No. 137. US Government Print Office.
44. Allmon, W.D., Harris, W.B. 2008. A new species of turritelline gastropod from a turritelline-dominated limestone in the Paleocene of North Carolina. *Journal of Paleontology*. 82: 442-446.

45. William J. Zinsmeister.1981.Middle to Late Eocene Invertebrate Fauna from the San Julian Formation at Punta Casamayor,Santa Cruz Province, Southern Argentina. Journal of Paleontology. 55: 1083-1102.