

Figure 1.

The accompanying figure presents a compilation of selected biostratigraphic data for the Late Cretaceous reference section near Pueblo Colorado; the sources for primary biostratigraphic data are indicated on the right side of the diagram by numbers keyed to the bibliographic information below. The figure also shows a lithostratigraphic correlation of the Pueblo section to the Angus and Portland core records, which is the basis for assignment of biozones to the core records. Lastly, the figure includes a summary of FAD levels for key taxa in the CL-1 core.

REFERENCES FOR SUPPLEMENTARY FIGURE 4

- 1 - COBBAN, W.A., and SCOTT, G.R., 1972, Stratigraphy and ammonite fauna of the Graneros Shale and Greenhorn Limestone near Pueblo, Colorado: United States Geological Survey Professional Paper v. P-0645, US Government Printing Office, 108 p.
- 2 - KAUFFMAN, E.G., 1985, Depositional history of the Graneros Shale (Cenomanian), Rock Canyon Anticline. In L. M. Pratt, E. G. Kauffman, and F. Zelt (eds.), Soc. Econ. Paleontologists, Mineralogists, Field Trip Guidebook No. 4; 1985 Midyear Meeting, Golden, CO p. 90-99.
- 3 - COBBAN, W.A., 1984, Mid-Cretaceous ammonite zones, Western Interior, United States. *Bull. Geol. Soc. Denmark*, v. 33: 71-89.
- 4 - COBBAN, W.A., 1985, Ammonite recored from Bridge Creek Member of Greenhorn Limestone at Pueblo Reservoirs State Recreation Area, Colorado, in Pratt, L.M., Kauffman, E.G., and Zelt, F.B., eds., *Fine-grained Deposits and Biofacies of the Cretaceous Western Interior Seaway: Evidence of Cyclic Sedimentary Processes*: SEPM, Field Trip Guidebook No. 4, p. 135-138.
- 5 - KAUFFMAN, E.G., SAGEMAN, B.B., KIRKLAND, J.I., ELDER, W.P., HARRIES, P.J., and VILLAMIL, T., 1993, Cretaceous molluscan biostratigraphy and biogeography, Western Interior Basin, North America. In: W.G.E. Caldwell and E.G. Kauffman (eds.), *Cretaceous Evolution of the Western Interior Basin, North America*, Special Publ. Geol. Assn. Canada 39: p. 397-434.
- 6 - SAGEMAN, B.B., 1985, High-resolution stratigraphy and paleobiology of the Hartland Shale Member: Analysis of an oxygen-deficient epicontinental sea. **In:** L.M. Pratt, E.G. Kauffman, and F.B. Zelt (eds.), *Fine-grained deposits and biofacies of the Cretaceous Western Interior Seaway: Evidence of cyclic sedimentary processes*. Society Economic Paleontologists Mineralogists, Field Trip Guidebook No. 4, Midyear Meeting, Aug. 12-14, 1985, Golden, Colorado, p. 110-121.
- 7 - SAGEMAN, B.B. and JOHNSON, C.C., 1985, Stratigraphy and paleobiology of the Lincoln Limestone Member, Greenhorn Limestone, Rock Canyon Anticline, Colorado. **In:** Pratt, L.M., Kauffman, E.G. and Zelt, F.B. (eds.), *Fine-grained deposits and biofacies of the Cretaceous Western Interior Seaway: Evidence of cyclic sedimentary processes*. Society Economic Paleontologists Mineralogists, Field Trip Guidebook No. 4, Midyear Meeting, Aug. 12-14, 1985, Golden, Colorado, p. 100-109.
- 8 - CARON, M., Dall'Agnolo, S., Accarie, H., Barrera, E., Kauffman, E.G., Amédro, F., and Robaszynski, F., 2006, High-resolution stratigraphy of the

Cenomanian/Turonian boundary interval at Pueblo (USA) and Wadi Bahloul (Tunisia): stable isotope and bio-events correlation, *Geobios* 39: p. 171-200.

- 9 - COBBAN, W.A., 1987, Some Middle Cenomanian (Upper Cretaceous) Acentrocerid Ammonites from the Western Interior of the United States, U.S.G.S. Professional Paper 1445, US Government Printing Office, 60 pp.
- 10 - FISHER, C. G., KAUFFMAN, E.G., WILHELM, and L. VON HOLT, 1985, The Niobrara transgressive hemicyclothem in central and eastern Colorado; the anatomy of a multiple unconformity, *SEPM Guidebook* 4, p. 184-198.
- 11 - GLENISTER, L., and KAUFFMAN, E. G., 1985, High resolution stratigraphy and depositional history of the Greenhorn Regressive Hemicyclothem, Rock Canyon Anticline, Pueblo, Colorado. In L. M. Pratt, E. G. Kauffman, and F. Zelt (eds.), *Soc. Econ. Paleontologists, Mineralogists, Field Trip Guidebook No. 4; 1985 Midyear Meeting*, Golden, CO. p. 170-183.
- 12 - SCOTT, G.R., and COBBAN, W.A., 1964, Stratigraphy of the Niobrara Formation at Pueblo, Colorado: U.S.G.S. Professional Paper, v. 454-L, US Government Printing Office, 30 pp., 11 plates.
- 13 - COBBAN, W.A., ERDMANN, C.E., LEMKE, R.W., and E.K. MAUGHN, 1976, Type sections of the Black Leaf and Marias River Formations (Cretaceous) of the Sweetgrass Arch, Montana. U.S.G.S. Professional Paper 974, US Government Printing Office, 65 pp.
- 14 - COBBAN, W.A., DYMAN, T.S., and PORTER, K.W., 2005, Paleontology and stratigraphy of upper Coniacian-middle Santonian ammonite zones and application to erosion surfaces and marine transgressive strata in Montana and Alberta: *Cretaceous Research*, v. 26, no. 3, p. 429-449.
- 15 - LANDMAN, N.H. and COBBAN, W.A., 2007, Redescription of the Late Cretaceous (late Santonian) ammonite *Desmoscaphites bassleri* Reeside, 1927, from the Western Interior of North America, *Rocky Mountain Geology*, v. 42, no. 2, 67-94.
- 16 - COBBAN, W.A., 1964, The Late Cretaceous cephalopod *Haresiceras* Reeside and its possible origin: U.S.G.S. Professional Paper 454-I, US Government Printing Office, 21 p., 3 pl.
- 17 - WALASZCZYK, I. and COBBAN, W.A., 1999, The Turonian-Coniacian boundary in the United States Western Interior: *Acta Geologica Polonica*, v. 48, p. 495-507.

_____ 2000, Inoceramid Faunas and Biostratigraphy of the Upper Turonian-Lower Coniacian of the Western Interior of the United States: Palaeontological Association [London] Special Paper, v. 64, 118 pp.

_____ 2006, Palaeontology and stratigraphy of the Middle-Upper Coniacian and Santonian inoceramids of the US Western Interior: *Acta Geologica Polonica*, v. 56, p. 241-348

_____ 2007, Inoceramid fauna and biostratigraphy of the upper Middle Coniacian-lower Middle Santonian of the Pueblo Section (SE Colorado, US Western Interior): *Cretaceous Research* v. 28, p. 132-142.

- 18 - GRIFI, M.D., PLINT, A.G., and WALASZCZYK, I., 2013, Rapidly changing styles of subsidence revealed by high-resolution mudstone allostratigraphy: Coniacian of Sweetgrass Arch area, southern Alberta and northern Montana, *Canadian Journal of Earth Sciences*, v. 50, p. 1–24 (2013) dx.doi.org/10.1139/cjes-2012-0031
- 19 - LOCKLAIR, R.E., and SAGEMAN, B.B., 2008, Cyclostratigraphy of the Upper Cretaceous Niobrara Formation, Western Interior, U.S.A: A Coniacian-Santonian orbital timescale: *Earth and Planetary Science Letters*, v. 269, p. 540-553.
- 20 - SAGEMAN, B.B., SINGER, B.S., MEYERS, S.R., WALASZCZYK, I., SEIWERT, S.E., CONDON, D.J., JICHA, B.R., OBRADOVICH, J.D., and SAWYER, D.A., 2014, in press, Integrating $^{40}\text{Ar}/^{39}\text{Ar}$, U-Pb, and astronomical clocks in the Cretaceous Niobrara Formation, Western Interior Basin, USA, *Geol. Soc. Amer. Bull.*
- 21- BALL, B.A., COBBAN, W.A., MEREWETHER, E.A., GRAUCH, R.I., MCKINNEY, K.C., and LIVO, K.E., 2010, Fossils, lithologies, and geophysical logs of the Mancos Shale from core hole USGS CL-1 in Montrose County, Colorado: U.S. Geological Survey Open-File Report 2009-1294, p. 38.