

Supplementary Data 1: Selected references for data used in Whidden and Jones (2012).

- ADAMS, C.G., 1970, A reconsideration of the East Indian Letter Classification of the Tertiary: Bulletin of the British Museum (Natural History), Geology, v. 19, p. 87-137.
- AFZAL, J., WILLIAMS, M. & ALDRIDGE, R.J., 2009, Revised stratigraphy of the lower Cenozoic succession of the Greater Indus Basin in Pakistan: Journal of Micropalaeontology, v. 28, p. 7-23.
- BANNER, F.T. & HODGKINSON, R.L., 1991, A revision of the foraminiferal subfamily Heterostegininae: Revista Espanola de Micropaleontologia, v. 23(2), p. 101-140.
- BOUDAGHER-FADEL, M.K., 2008, *Evolution and geological significance of larger benthic foraminifera*. Elsevier, Amsterdam (Developments in Palaeontology and Stratigraphy, No. 21).
- BOUDAGHER, M.K. & BANER, F.T., 1999, Revision of the stratigraphic significance of the Oligocene-Miocene "Letter Stages": Revue de Micropaleontologie, v. 42(2), p.93-97.
- BOUDAGHER-FADEL, M.K. & PRICE, G.D., 2010, Evolution and paleogeographic distribution of the lepidocyclinids: Journal of Foraminiferal Research, v. 40(1), p. 79-108.
- CAHUZAC, B. & POIGNANT, A., 1997, Essai de biozonation de l'Oligo-Miocene dans less bassins europeens a l'aide des grands foraminifères neritiques: Bulletin de la Societe Geologique de la France, v. 168(2): 155-169.
- CAUDRI, C.M.B., 1996. The larger foraminifera of Trinidad (West Indies). *Eclogae Geologicae Helvetiae*, 89(3), p. 1137-1309.
- CHAPRONIERE, G.C.H., 1981, Australasian mid-Tertiary larger foraminiferal associations and their bearing on the East Indian Letter Classification: Bureau of Mineral Resources Journal of Australian Geology and Geophysics, v. 6, p. 145-151.
- EAMES, F.E., 1952, A contribution to the study of the Eocene of western Pakistan and western India: Quarterly Journal of the Geological Society, v. 107, p. 159-200.
- FLANDRIN, J., 1938, Contribution a l'étude paleontologique du Nummulitique algérien: Materieux pour la Carte Geologique de l'Algérie (1e Serie, Paleontologie, No. 8).
- HENSON, F.R.S., 1948 Larger imperforate foraminifera from South Western Asia: British Museum (Natural History).
- HOTTINGER, L., 1960, Über Paleocaene und Eocaene Alveolinen: Eclogae Geologicae Helvetiae, v. 53, p. 265-283.
- HOTTINGER, L., 1977, Foraminifères operculiniformes: Memoires de la Musee National d'Histoire Naturelle, C, v. 40, 159pp.
- HOTTINGER, L. & DROBNE, K., 1988, Alveolines Tertiaires: quelques problems lies a la conception de l'espèce: Revue de Paleobiologie, Volume Special, v. 2, p. 65-685.

- JONES, R.W., 1997 Aspects of the Cenozoic stratigraphy of the northern Sulaiman Ranges, Pakistan: *Journal of Micropalaeontology*, v. 16, p. 51-58.
- JONES, R.W., 1999, Marine Invertebrate (Chiefly Foraminiferal) Evidence for the Palaeogeography of the Oligocene-Miocene of Western Eurasia, and Consequences for Terrestrial Vertebrate Migration: *in* Agusti, J., Rook, L. & Andrews, P., eds.,: Hominoid evolution and climatic change in Eurasia. Vol. 1: Climatic and environmental change in the Neogene of Europe, Cambridge University Press, p. 274-308.
- JONES, R.W. & RACEY, A., 1994, Cenozoic Stratigraphy of the Arabian Peninsula and Gulf: *in* Simmons, M.D. (ed.), *Micropalaeontology and Hydrocarbon Exploration in the Middle East*, London; Chapman & Hall, p. 273-307.
- KURESHY, A.A., 1978, Tertiary larger foraminiferal zones of Pakistan: *Revista Espanola de Micropaleontologia*, v. 10(3), p. 467-483.
- LESS, G., 1987, Paleontology and stratigraphy of the European Orthophragmidae: *Geologia Hungarica (Palaeontology)*, v. 51, 373pp.
- MELLO E SOUSA, S.H., FAIRCHILD, T.R. & TIBANA, P., 2003, Cenozoic biostratigraphy of larger foraminifera from the Foz do Amazonas Basin, Brazil: *Micropaleontology*, v. 49, p. 253-266.
- MORLEY DAVIES, A. & EAMES, F.E., 1971, *Tertiary Faunas*: London; George Allen & Unwin.
- RACEY, A., 1994, Biostratigraphy and palaeobiogeographic significance of Tertiary nummulitids (Foraminifera) from northern Oman: *in* Simmons, M.D., ed., *Micropalaeontology and Hydrocarbon Exploration in the Middle East*, London; Chapman & Hall, p. 343-370.
- RACEY, A., 1995, Lithostratigraphy and larger foraminiferal (nummulitid) biostratigraphy of the Tertiary of northern Oman: *Micropaleontology*, v. 41(Supplement), p. 1-123.
- RAJU, D.S.N., 1974, Study of Indian Miogypsinidae: *Utrecht Micropalaeontological Bulletin*, v. 9, p. 1-148.
- ROBINSON, E., 1968. Stratigraphic ranges of some larger benthonic foraminifera in Jamaica: *Transactions of the Fourth Caribbean Geological Conference*, Trinidad, 1965, p. 189-194.
- SAMANTA, B.K., 1969, Taxonomy and stratigraphy of the Indian species of *Discocyclina*: *Geol. Mag.*, v. 106(2), p. 115-129.
- SCHAUB, H., 1981. Nummulites et Assilines de la Tethys Paleogene. Taxinomie, phylogénèse et biostratigraphie: *Schweizerische Palaontologisches Abhandlungen*, v. 104-106, 236 pp.
- SEIGLIE, G.A. & FROST, S.H., 1979. Significance of middle Tertiary large foraminifera common to West Africa and Caribbean: *American Association of Petroleum Geologists Bulletin*, v. 63, p. 525.
- SERRA-KIEL, J., HOTTINGER, L., CAUS, E. ET AL., 1998, Larger foraminiferal biostratigraphy of the Tethyan Palaeocene and Eocene: *Bulletin de la Societe Geologique de la France*, v. 169(2), p. 281-299.

SMOUT, A.H., 1954, Lower Tertiary foraminifera of the Qatar Peninsula: British Museum (Natural History).

VAN VESSEM, E.J., 1978, Study of Lepidocylinidae from Southeast Asia, particularly from Java and Borneo: Utrecht Micropalaeontological Bulletin, v. 19, p. 1-163.

WHITE, M.R., 1992, On species identification in the foraminiferal genus *Alveolina* (Late Paleocene-Middle Eocene): Journal of Foraminiferal Research, v. 22(1), p. 52-70.

WIELANDT, U., 1996, Benthic foraminiferal paleoecology and microfacies investigations of Paleogene sediments from the Farafra Oasis, Western Desert, Egypt: Tubinger Mikropalaontologische Mitteilungen, v. 13, p. 1-78.